

Published in final edited form as:

*Psychol Addict Behav.* 2016 February ; 30(1): 101–105. doi:10.1037/adb0000112.

## The Effects of E-Cigarette Visual Appearance on Craving and Withdrawal Symptoms in Abstinent Smokers

Lynne Dawkins, PhD<sup>a</sup>, Marcus Munafò, PhD<sup>b</sup>, Gina Christoforou, BSc<sup>a</sup>, Naomi Olumegbon, BSc<sup>a</sup>, and Kirstie Soar, PhD<sup>a</sup>

<sup>a</sup>School of Psychology, University of East London, Water Lane, Stratford, London, E15 4LZ

<sup>b</sup>MRC Integrative Epidemiology Unit, UK Centre for Tobacco and Alcohol Studies, and School of Experimental Psychology, University of Bristol, 12a Priory Road, Bristol, BS8 1TU

### Abstract

**Introduction**—Electronic cigarette (e-cigarette) use is becoming increasingly popular among smokers and there is a plethora of devices available. Nicotine delivery is clearly important for reducing tobacco craving and withdrawal symptoms, but other sensor-motor aspects of e-cigarettes (such as visual appearance) may contribute to this effect. This study explored whether it is important for an e-cigarette to visually resemble a tobacco cigarette in order to reduce craving and withdrawal symptoms.

**Methods**—Sixty-three abstinent smokers (40% female, aged 18–65 years) who were not current e-cigarette users were randomly allocated to take ten 3-second puffs from either a white or a red first generation e-cigarette. Current craving (urge to smoke) and nicotine withdrawal symptoms (using the Mood and Physical Symptoms Scale; MPSS) were measured before and ten minutes after use.

**Results**—Linear regression revealed higher craving and withdrawal symptoms in the red versus the white condition but only among those who were e-cigarette naive (craving:  $B = .76$ ,  $p = .009$ ; withdrawal symptoms:  $B = 2.18$ ,  $p = 0.009$ ), not among those with e-cigarette experience (craving:  $B = -.08$ ,  $p = 0.89$ ; withdrawal symptoms:  $B = .24$ ,  $p = .81$ ), and these effects differed between groups ( $p = 0.04$  and  $0.01$  for craving and withdrawal symptoms respectively).

**Conclusion**—Cigarette-like appearance was associated with a greater reduction in craving and withdrawal symptoms but only for those with no prior e-cigarette experience. This effect, putatively mediated via classical conditioning or expectancies, may aid understanding of smokers' initial preferences for 'cigalike' e-cigarette devices.

---

Corresponding author: Lynne Dawkins; L.E.Dawkins@uel.ac.uk; Tel: +44 (0)20 8223 4421; Fax: +44 (0)20 8223 4937.

#### Declaration of Interests

LD has conducted research for electronic cigarette companies for which the University of East London has received funding. All other authors declare no conflict of interest.

MM is a member of the UK Centre for Tobacco and Alcohol Studies, a UKCRC Public Health Research: Centre of Excellence. Funding from British Heart Foundation, Cancer Research UK, Economic and Social Research Council, Medical Research Council, and the National Institute for Health Research, under the auspices of the UK Clinical Research Collaboration, is gratefully acknowledged.

## Keywords

Electronic cigarettes; craving; withdrawal symptoms; sensori-motor aspects; visual appearance; cigalike

---

## 1 Introduction

Electronic cigarettes (e-cigarettes) are battery-powered devices that deliver nicotine via an aerosol that is inhaled. Recent evidence supports their efficacy as a smoking cessation aid with quit rates at least equivalent to Nicotine Replacement Therapy (NRT; McRobbie et al., 2014; Brown et al., 2014). Although a relatively new phenomenon, public awareness and regular use of e-cigarettes has increased rapidly, doubling between 2010 and 2011 in the US (King et al., 2013) and increasing from 3% to 18% between in 2010 and 2014 in Great Britain (Action on Smoking and Health, [ASH], 2014) although use in England has dipped recently (West & Brown, 2015).

E-cigarette use resembles the act of smoking: the user holds the device and draws on it like a cigarette; the aerosol produced is drawn into the lungs and exhaled like smoke; tobacco (or menthol) flavouring mimics the taste of inhaled tobacco smoke and many (first generation) device look exactly like tobacco cigarettes (commonly known as ‘cigalikes’). Although regular e-cigarette users tend to use devices which deviate from a cigarette-like appearance (second or third generation devices; Dawkins et al., 2013), ‘cigalikes’ currently account for 65% of the e-cigarette market share (Herzog, Gerberi & Scott, 2014) and are more commonly found in retail outlets across the UK and US. Such devices may appeal to new e-cigarette users; in a recent study of 100 (e-cigarette naïve) smokers who were asked to choose between a first or second generation device, 50% opted for a first generation ‘cigalike’ and did so because it looked like a tobacco cigarette (Dawkins et al., 2014).

Although nicotine is clearly a critical component of tobacco dependence, a growing body of evidence points to the role of non-nicotine, sensorimotor factors in supporting smoking behaviour. Anecdotally, smokers prefer smoking to other forms of nicotine administration (e.g., patch, gum, nasal spray), possibly due to the rapid nicotine delivery although the sensory and tactile components of smoking, including the hand-mouth activity, taste, smell and sensations in the respiratory tract are all strongly endorsed (Parrott & Craig, 1995). Smokers have also been shown to prefer smoking a denicotinised cigarette over receiving intravenous nicotine (Rose, et al., 2010) and denicotinised tobacco smoking can alleviate nicotine withdrawal symptoms and craving (Barrett, 2010; Perkins et al., 2010; Rose et al., 2010). These effects likely arise as a consequence of sensorimotor cues (e.g., visual appearance, hand-mouth action, taste and smell of smoke) imbuing some of the rewarding properties of smoking given their repeated pairing with nicotine administration via classical conditioning.

Acute e-cigarette use can also reduce tobacco craving and withdrawal symptoms in abstinent smokers (Bullen et al. 2010; Dawkins, Turner & Crowe, 2013; Vansickel et al., 2010) and non-nicotine placebo devices have also been shown to be effective (Dawkins et al., 2012). Nevertheless, the importance of the visual similarity to tobacco cigarettes has not been

systematically explored. We have recently reported that a first and second generation device were equally effective in suppressing tobacco craving and withdrawal symptoms after acute use (Dawkins et al., 2014). Although not measured in this study, this may have been due to similar nicotine absorption, or because the cigarette-like appearance of the first generation device elicited a conditioned drug-like response (Stewart, de Wit & Eikelboom, 1984; Carter & Tiffany, 2001) that made up for inferior nicotine delivery in first-generation devices (Farsalinos et al., 2014). Given the documented importance of sensorimotor aspects of smoking, it is possible that an e-cigarette that looks visually similar to a tobacco cigarette may be more useful for reducing acute cigarette craving and withdrawal symptoms than one that does not. Here we tested the importance of visual appearance using a first generation 'cigalike' device and varied just one aspect of visual appearance: colour (white: visually similar to a cigarette vs. red: visually dissimilar to a cigarette), with nicotine content and all other components remaining identical (orange cartridge, size, shape, flavour). We hypothesised that, relative to the red condition, abstinent smokers in the white condition, would show a greater reduction in craving and withdrawal symptoms following a standard 10 puff regime.

## 2 Method

### 2.1 Participants

Sixty-three abstinent smokers (27 [40%] female,) aged between 18 and 65 years (mean 27, SD 11) were recruited via poster advertisements at the University of East London (UEL), e-mail and social networking websites. All were daily smokers who had been smoking for at least one year, not currently interested in quitting and had not used an e-cigarette in the past 30 days.

### 2.2 Procedure and Measures

Following informed consent, participants provided a breath carbon monoxide (CO) sample measured using a Bedfont piCO+ smokerlyser to confirm overnight abstinence (>10 hours) with a level less than 10 ppm required for participation. Baseline demographic information, previous experience with e-cigarettes and smoking behaviour including severity of nicotine/tobacco dependence measured using the 6-item Fagerström Test for Cigarette Dependence (FTCD; Fagerström, 2012) was collected. Participants then rated their current urge to smoke (craving) on a single-item 5-point rating scale, from 0 (no urges) to 5 (extremely strong urge) and nicotine withdrawal symptoms were assessed using the Mood and Physical Symptoms Scale (MPSS; West & Hajek, 2004) which includes six core items (depressed mood, anxiousness, irritability, restlessness, hunger, poor concentration) rated on a scale from 1 (not at all) to 5 (extremely). Following random allocation (via coin toss) to group (red vs. white), all participants took ten, three-second puffs on a regular strength (18 mg/ml) 'tobacco' flavour rechargeable e-cigarette from E-lites with a 30-second inter-puff interval (consistent with that of Vansickle & Eissenberg, 2012). After ten minutes, current strength of urge to smoke and withdrawal symptoms were rated for a second time.

## 2.3 Statistical Analysis

Linear regression was used to assess the effect of condition (red vs. white) on craving and withdrawal symptom scores. Primary analyses were conducted adjusted for baseline craving/MPSS, age, sex, ethnicity and cigarettes smoked per day (CPD). Given that some participants reported prior use of an e-cigarette (current users were excluded), post hoc analyses tested for an interaction between condition and prior e-cigarette use. Where evidence for an interaction effect was observed, secondary analyses stratified by prior e-cigarette use were used to explore the nature of the interaction. Our sample size provided 80% power to detect an effect size of  $d = 0.72$  at an alpha level of 5%, equivalent to a difference of 0.5 points on the craving measure (assuming  $SD = 1$ ) and 1 point on the MPSS (assuming  $SD = 3$ ). All analyses were conducted with IBM SPSS Statistics (version 20).

## 3 Results

The two groups did not differ on any demographic or smoking related variables (see Table 1). Twenty-two participants had previously used an e-cigarette. Prior e-cigarette use ranged from 1 to 20 occasions (mean mean 3;  $SD$  4) with 68% reporting using only once or twice.

### 3.1 Craving

In the primary analysis, we observed no strong evidence of a main effect of condition on craving ( $B = +0.376$ , 95% CI  $-0.117$  to  $+0.870$ ,  $p = 0.132$ ). However, there was evidence of a condition  $\times$  prior e-cigarette use interaction ( $B = -0.012$ , 95% CI  $-0.024$  to  $-0.001$ ,  $p = 0.038$ ). Secondary analyses stratified by prior e-cigarette use confirmed higher craving in the red compared with the white condition among those who had not used an e-cigarette in the past ( $B = +0.775$ , 95% CI  $+0.205$  to  $+1.344$ ,  $p = 0.009$ ), but not among those who had ( $B = -0.079$ , 95% CI  $-1.283$  to  $+1.126$ ,  $p = 0.891$ ). These results are shown in Table 2.

### 3.2 Withdrawal Symptoms

In the primary analysis, we observed higher withdrawal symptom scores in the red compared with the white condition ( $B = +1.887$ , 95% CI  $+0.606$  to  $+3.169$ ,  $p = 0.005$ ). There was also evidence of a condition  $\times$  prior e-cigarette use interaction ( $B = -0.038$ , 95% CI  $-0.067$  to  $-0.009$ ,  $p = 0.012$ ). Secondary analyses stratified by prior e-cigarette use confirmed higher withdrawal symptoms in the red compared with the white condition among those who had not used an e-cigarette in the past ( $B = +2.177$ , 95% CI  $+0.573$  to  $+3.781$ ,  $p = 0.009$ ) but not for those who had ( $B = +0.243$ , 95% CI  $-1.850$  to  $+2.337$ ,  $p = 0.808$ ). These results are shown in Table 2.

## 4 Discussion

Although many smokers choosing an e-cigarette for the first time opt for 'cigalike' devices (Dawkins et al., 2014), whether the visual similarity of an e-cigarette to a tobacco cigarette is important for the alleviation of craving and withdrawal symptoms has not been explored. In abstinent smokers who had not used an e-cigarette in the past, we observed a greater reduction in craving and withdrawal symptoms after using the white (visually similar to a cigarette) compared with the red (visually dissimilar to a cigarette) e-cigarette. For

participants with previous e-cigarette experience, no such difference was found, suggesting that cigarette-like appearance may be an important factor in alleviating acute craving and withdrawal symptoms but only for e-cigarette naive smokers. However, this moderating effect of prior experience was not specifically hypothesised, and therefore should be treated with caution until replicated independently.

Sensorimotor cues such as visual appearance act as conditioned reinforcers that can become moderately reinforcing in their own right (Stewart, de Wit & Eikelboom, 1984). Given that nicotine content was identical in the visually similar and dissimilar conditions, a conditioned drug-like response is consistent with the greater reduction in craving/withdrawal symptoms in the visually similar condition observed here. An alternative explanation is that the visually similar (white) e-cigarette was perceived as more effective, although unfortunately expectations regarding effectiveness were not measured. Either way, whether via an unconscious conditioning mechanism or a conscious expectancy effect or indeed both, there is evidence here that acute craving and withdrawal symptom alleviation is not exclusively attributable to nicotine delivery.

In the current study, the effect of visual appearance was only found in participants with no e-cigarette experience and most of those who had previously tried an e-cigarette had done so only once or twice (68%). Whilst visual similarity to a cigarette may therefore be useful for reducing craving and withdrawal symptoms in the initial stages of e-cigarette use, this effect may dissipate fairly quickly. This may explain why many smokers initially opt for 'cigalikes' (Dawkins et al., 2014) whilst continuing users transition to cigarette dissimilar second and third generation devices (Dawkins et al., 2013; Etter & Bullen, 2011; Farsalinos et al., 2013; McQueen, Tower & Sumner, 2011) which may offer superior nicotine delivery (Farsalinos et al., 2014).

Limitations to the present study include the exclusive focus on visual cigarette-like appearance. Whilst this makes for a more controlled experimental study, it does not acknowledge the importance of other sensorimotor factors such as vapour production, taste and design. Moreover, red 'cigalike' devices are less commonly available and it is the second and third generation refillable devices that are competing with 'cigalikes' in the marketplace. We have previously compared a 'cigalike' and second generation device and found they were equally effective at reducing craving and withdrawal symptoms (Dawkins et al., 2014) but differences in nicotine delivery between the devices were not measured and complicates interpretation. The clinical significance of the findings for longer-term smoking abstinence also remain unclear given the effects were limited to craving and withdrawal symptom reduction in a sub-sample of e-cigarette naive smokers after only 10 e-cigarette puffs. A longer-term follow up study would determine how long the effect is retained and whether it has any predictive utility for smoking cessation.

In conclusion, compared with a cigarette-dissimilar device, an e-cigarette that visually resembled a tobacco cigarette was associated with a greater reduction of tobacco craving and withdrawal symptoms in e-cigarette naive abstinent smokers. Similar effects were not observed in those with previous e-cigarette experience suggesting the effect may be short-lived. This may be one of the reasons why many smokers favour 'cigalikes' initially but

often transition to cigarette-dissimilar second and third generation devices over time (Dawkins et al., 2013; 2014, McQueen, Tower & Sumner, 2011).

## Acknowledgements

No funding was provided for this study although thanks are due to E-Lites for supplying the e-cigarettes. The company had no involvement in the design, conduct or write up of the study.

## References

- ASH. Ash Fact Sheet: Use of electronic cigarettes in Great Britain, April 2014. 2014. [http://www.ash.org.uk/files/documents/ASH\\_891.pdf](http://www.ash.org.uk/files/documents/ASH_891.pdf)
- Barrett SP. The effects of nicotine, denicotinized tobacco, and nicotine-containing tobacco on cigarette craving, withdrawal, and self-administration in male and female smokers. *Behavioral Pharmacology*. 2010; 21(2):144–152. DOI: 10.1097/FBP.0b013e328337be68
- Brown J, Beard E, Kotz D, Michie S, West R. Real-world effectiveness of e-cigarettes when used to aid smoking cessation: a cross-sectional population study. *Addiction*. 2014; 109:1531–40. DOI: 10.1111/add.12623 [PubMed: 24846453]
- Bullen C, McRobbie H, Thornley S, Glover M, Lin R, Laugesen M. Effects of an electronic nicotine delivery device (e cigarette) on desire to smoke and withdrawal, user preferences and nicotine delivery: randomised cross-over trial. *Tobacco Control*. 2010; 19:98–103. DOI: 10.1136/tc.2009.031567 [PubMed: 20378585]
- Carter BL, Tiffany ST. The cue-availability paradigm: the effects of cigarette availability on cue reactivity in smokers. *Experimental and Clinical Psychopharmacology*. 2001; 9(2):183–90. [PubMed: 11518094]
- Dawkins L, Kimber C, Puwanesarasa Y, Soar K. First- versus second-generation electronic cigarettes: predictors of choice and effects on urge to smoke and withdrawal symptoms. *Addiction*. 2014; published online 22 Dec.2014. doi: 10.1111/add.12807
- Dawkins L, Turner J, Crowe E. Nicotine derived from the electronic cigarette improves time-based prospective memory in abstinent smokers. *Psychopharmacology*. 2013; 227:377–384. DOI: 10.1007/s00213-013-2983-2 [PubMed: 23344557]
- Dawkins L, Turner J, Hasna S, Soar K. The electronic-cigarette: Effects on desire to smoke, withdrawal symptoms and cognition. *Addictive Behaviors*. 2012; 37:970–973. DOI: 10.1016/j.addbeh.2012.03.004 [PubMed: 22503574]
- Dawkins L, Turner J, Roberts A, Soar K. ‘Vaping’ profiles and preference: an online survey of electronic cigarette users. *Addiction*. 2013; 108(6):1115–1125. DOI: 10.1111/add.12150 [PubMed: 23551515]
- Etter J, Bullen C. Electronic cigarette: users profile, utilization, satisfaction and perceived efficacy. *Addiction*. 2011; 106:2017–2028. DOI: 10.1111/j.1360-0443.2011.03505.x [PubMed: 21592253]
- Farsalinos KE, Romagna G, Tsiapras D, Kyrozoopoulos S, Voudris V. Evaluating nicotine levels selection and patterns of electronic cigarette use in a group of ‘vapers’ who achieved complete substitution of smoking. *Substance Abuse: Research and Treatment*. 2013; 7:139–146. DOI: 10.4137/SART.S12756 [PubMed: 24049448]
- Farsalinos KE, Spyrou A, Tsimopoulou K, Stefanopoulos C, Romagna G, Voudris V. Nicotine absorption from electronic cigarette use: comparison between first and new-generation devices. *Scientific Reports*. 2014; 4:4133.doi: 10.1038/srep04133 [PubMed: 24569565]
- Fagerström KO. Determinants of tobacco use and renaming the FTND to the Fagerström Test for Cigarette Dependence. *Nicotine and Tobacco Research*. 2012; 14:75–78. DOI: 10.1093/ntr/ntr137 [PubMed: 22025545]
- Herzog, B.; Gerberi, J.; Scott, A. Equity Research – Tobacco: Vapor World Expo – Key Takeaways. Wells Fargo Securities; 2014. May 12, 2014
- King BA, Alam S, Promoff G, Arrazola R, Dube SR. Awareness and ever use of electronic cigarettes among US adults. *Nicotine and Tobacco Research*. 2013; 15:1623–1627. DOI: 10.1093/ntr/ntt013 [PubMed: 23449421]

- McQueen A, Tower S, Sumner W. Interviews with 'vapers': implications for future research with electronic cigarettes. *Nicotine and Tobacco Research*. 2011; 13(9):860–7. DOI: 10.1093/ntr/ntr088 [PubMed: 21571692]
- McRobbie H, Bullen C, Hartmann-Boyce J, Hajek P. Electronic cigarettes for smoking cessation and reduction. *Cochrane Database of Systematic Reviews*. 2014; (12) Art. No.: CD010216. doi: 10.1002/14651858.CD010216.pub2
- Parrott AC, Craig D. Psychological functions served by nicotine chewing gum. *Addictive Behaviors*. 1995; 20(3):271–278. [PubMed: 7653311]
- Perkins KA, Karelitz JL, Conklin CA, Sayette MA, Giedgowd GE. Acute negative affect relief from smoking depends on the affect situation and measure but not on nicotine. *Biological Psychiatry*. 2010; 67(8):707–714. DOI: 10.1016/j.biopsych.2009.12.017 [PubMed: 20132927]
- Rose JE, Salley A, Behm FM, Bates JE, Westman EC. Reinforcing effects of nicotine and non-nicotine components of cigarette smoke. *Psychopharmacology*. 2010; 210:1–12. DOI: 10.1007/s00213-010-1810-2
- Stewart J, de Wit H, Eikelboom R. Role of unconditioned and conditioned drug effects in the self-administration of opiates and stimulants. *Psychological Review*. 1984; 91:251–268. [PubMed: 6571424]
- Vansickel AR, Cobb CO, Weaver MF, Eissenberg TE. A clinical laboratory model for evaluating the acute effects of electronic 'cigarettes': Nicotine delivery profile and cardiovascular and subjective effects. *Cancer Epidemiology, Biomarkers & Prevention*. 2010; 19:1945–1953. DOI: 10.1158/1055-9965.EPI-10-0288
- West, R.; Brown, J. Latest trends on smoking in England from the Smoking Toolkit Survey. 2015. <http://www.smokinginengland.info/latest-statistics/>
- West R, Hajek P. Evaluation of the mood and physical symptoms scale (MPSS) to assess cigarette withdrawal. *Psychopharmacology*. 2004; 177:195–199. [PubMed: 15179542]

**Table 1**

Demographic and smoking related information for the visually similar (white) vs. visually dissimilar (red) e-cigarette groups

	<b>WHITE N = 33</b>	<b>RED N = 30</b>
	<b>Mean (SD)</b>	<b>Mean (SD)</b>
Age	25.48 (9.27)	27.83 (2.21)
CPD	12.58 (8.05)	11.43 (5.42)
Years Smoked	8.92 (7.24)	7.98 (5.23)
FTND	5.45 (2.69)	5.13 (1.87)
	<b>N (%)</b>	<b>N (%)</b>
Female	11 (33%)	16 (53%)
Previous e-cigarette use (Yes)	9 (27%)	13 (43%)
European ethnicity	16 (48%)	20 (67%)

CPD: cigarettes per day; FTND: Fagerstrom Test of Nicotine Dependence

**Table 2**

Mean (SD) Craving (urge to smoke) and Withdrawal Symptoms (MPSS scores) at baseline and ten minutes after e-cigarette use for the visually similar (white) vs. visually dissimilar (red) condition

	WHITE		RED	
	Baseline Mean (SD)	Post E-cig Use Mean (SD)	Baseline Mean (SD)	Post E-cig Use Mean (SD)
<b>Urge to smoke</b>	2.70 (1.29)	1.76 (1.20)	2.63 (0.96)	1.93 (1.05)
All participants	2.67 (1.34)	1.75 (1.15)	2.35 (0.86)	2.18 (0.88)
No prior use	2.78 (1.20)	1.78 (1.39)	3.00 (1.00)	1.62 (1.19)
Prior use				
<b>MPSS</b>	14.52 (4.40)	10.52 (3.20)	13.30 (3.90)	11.30 (3.97)
All participants	14.33 (4.34)	10.75 (3.53)	14.24 (3.77)	12.76 (4.16)
No prior use	15.00 (4.77)	9.89 (2.15)	12.08 (3.86)	9.38 (2.81)
Prior use				